

SANTA CRUZ COUNTY'S HEALTH STATUS PROFILE FOR 2000

| MORTALITY | | | | | | | | |
|------------|-------------------------|----------------------------|------------------|-------------------------|-----------------------------------|-------|-----------------------------------|--------------------|
| RANK ORDER | HEALTH STATUS INDICATOR | 1996-1998 DEATHS (AVERAGE) | CRUDE DEATH RATE | AGE-ADJUSTED DEATH RATE | 95% CONFIDENCE LIMITS LOWER UPPER | | STATEWIDE AGE-ADJUSTED DEATH RATE | NATIONAL OBJECTIVE |
| 9 | ALL CAUSES | 1,671.0 | 675.9 | 375.8 | 354.7 | 397.0 | 425.7 | N/E |
| 15 | MOTOR VEHICLE ACCIDENTS | 27.3 | 11.1 | 10.8 | 6.5 | 15.1 | 11.4 | 14.2 |
| 12 | UNINTENTIONAL INJURIES | 66.3 | 26.8 | 23.6 | 17.5 | 29.8 | 24.2 | 29.3 |
| 7 | FIREARM INJURIES | 17.3 | 7.0 * | 6.5 * | 3.2 | 9.8 | 11.6 | 11.6 |
| 19 | HOMICIDE | 8.7 | 3.5 * | 3.7 * | 1.1 | 6.3 | 9.0 | 7.2 |
| 26 | SUICIDE | 27.3 | 11.1 | 10.0 | 6.0 | 14.0 | 9.4 | 10.5 |
| 6 | ALL CANCERS | 364.3 | 147.4 | 97.0 | 86.0 | 108.1 | 110.3 | 130.0 |
| 2 | LUNG CANCER | 80.3 | 32.5 | 22.8 | 17.3 | 28.2 | 30.0 | 42.0 |
| 40 | FEMALE BREAST CANCER | 35.7 | 28.8 | 19.7 | 12.7 | 26.8 | 18.3 | 20.6 |
| 6 | CORONARY HEART DISEASE | 375.0 | 151.7 | 68.1 | 59.7 | 76.5 | 93.9 | 100.0 |
| 12 | CEREBROVASCULAR DISEASE | 129.0 | 52.2 | 21.7 | 17.2 | 26.2 | 25.3 | 20.0 |
| 30 | DRUG-RELATED DEATHS | 19.7 | 8.0 | 6.9 * | 3.8 | 10.0 | 7.5 | 3.0 |

| MORBIDITY | | | | | | | | |
|------------|-------------------------|---------------------------|-----------------|-----------------------------------|-------|---------------------------|--------------------|--|
| RANK ORDER | HEALTH STATUS INDICATOR | 1996-1998 CASES (AVERAGE) | CRUDE CASE RATE | 95% CONFIDENCE LIMITS LOWER UPPER | | STATEWIDE CRUDE CASE RATE | NATIONAL OBJECTIVE | |
| 33 | AIDS INCIDENCE | 19.33 | 7.82 | 4.33 | 11.31 | 17.31 | 43.00 | |
| 52 | MEASLES INCIDENCE | 0.67 | 0.27 * | 0.00 | 0.92 | 0.08 | 0.00 | |
| 28 | TUBERCULOSIS INCIDENCE | 15.00 | 6.07 * | 3.00 | 9.14 | 12.37 | 3.50 | |
| 38 | SYPHILIS INCIDENCE | 0.67 | 0.27 * | 0.00 | 0.92 | 1.24 | 4.00 | |

| INFANT DEATH | | | | | | | | |
|--------------|----------------------------|----------------------------|--------------------------------|-----------------------------------|------|--|--------------------|--|
| RANK ORDER | HEALTH STATUS INDICATOR | 1994-1996 DEATHS (AVERAGE) | BIRTH COHORT INFANT DEATH RATE | 95% CONFIDENCE LIMITS LOWER UPPER | | STATEWIDE BIRTH COHORT INFANT DEATH RATE | NATIONAL OBJECTIVE | |
| 19 | INFANT DEATHS: ALL RACES | 19.7 | 5.6 | 3.1 | 8.0 | 6.4 | 7.0 | |
| 21 | INFANT DEATHS: ASIAN/OTHER | 0.3 | 2.9 * | 0.0 | 12.7 | 5.3 | N/E | |
| 4 | INFANT DEATHS: BLACK | 0.0 | 0.0 + | - | - | 13.7 | 11.0 | |
| 32 | INFANT DEATHS: HISPANIC | 10.0 | 5.9 * | 2.3 | 9.6 | 6.0 | N/E | |
| 23 | INFANT DEATHS: WHITE | 9.3 | 5.5 * | 2.0 | 9.0 | 5.9 | N/E | |

| NATALITY | | | | | | | | |
|------------|-----------------------------|----------------------------|---------|-----------------------------------|------|-------------------|--------------------|--|
| RANK ORDER | HEALTH STATUS INDICATOR | 1996-1998 BIRTHS (AVERAGE) | PERCENT | 95% CONFIDENCE LIMITS LOWER UPPER | | STATEWIDE PERCENT | NATIONAL OBJECTIVE | |
| 14 | LOW BIRTHWEIGHT INFANTS | 172.7 | 5.0 | 4.2 | 5.7 | 6.1 | 5.0 | |
| 16 | LATE OR NO PRENATAL CARE | 564.7 | 16.5 | 15.2 | 17.9 | 18.4 | 10.0 | |
| 31 | ADEQUATE/ADEQUATE PLUS CARE | 2,241.7 | 67.0 | 64.2 | 69.8 | 70.5 | 90.0 ^ | |

| RANK ORDER | HEALTH STATUS INDICATOR | 1996-1998 BIRTHS (AVERAGE) | AGE-SPECIFIC BIRTH RATE | 95% CONFIDENCE LIMITS LOWER UPPER | | STATEWIDE AGE-SPECIFIC BIRTH RATE | NATIONAL OBJECTIVE |
|------------|------------------------------|----------------------------|-------------------------|-----------------------------------|------|-----------------------------------|--------------------|
| 23 | BIRTHS TO MOTHERS AGED 15-19 | 353.3 | 42.9 | 38.4 | 47.4 | 57.2 | N/E |

| BREASTFEEDING | | | | | | | | |
|---------------|--------------------------|------------------|---------|-----------------------------------|------|-------------------|--------------------|--|
| RANK ORDER | HEALTH STATUS INDICATOR | BIRTHS (AVERAGE) | PERCENT | 95% CONFIDENCE LIMITS LOWER UPPER | | STATEWIDE PERCENT | NATIONAL OBJECTIVE | |
| 1 | BREASTFEEDING INITIATION | 3,171.3 | 92.4 | 89.2 | 95.6 | 78.4 | 75.0 | |

| CENSUS | | | | | | | | |
|------------|-----------------------------|-------------|---------|-----------------------------------|------|-------------------|--------------------|--|
| RANK ORDER | HEALTH STATUS INDICATOR | 1990 NUMBER | PERCENT | 95% CONFIDENCE LIMITS LOWER UPPER | | STATEWIDE PERCENT | NATIONAL OBJECTIVE | |
| 15 | PERSONS UNDER 18 IN POVERTY | 6,280 | 11.9 | 11.6 | 12.2 | 18.2 | N/E | |

N/E: National Objective for the Year 2000 has not been established.

^ Year 2010 National Health Objective.

* Rate or percent unreliable; relative standard error greater than or equal to 23%.

+ Rate or percent indeterminate; no (zero) events.

- Upper and lower limits at the 95% confidence level are not calculated for no (zero) events.

Note: Crude death rates, crude case rates, and age-adjusted death rates are per 100,000 population. Birth cohort infant death rates are per 1,000 live births. Age-specific birth rates are per 1,000 population.

Sources: Department of Health Services: Center for Health Statistics, Birth and Death Statistical Master Files, 1996-1998, and Birth Cohort Files, 1994-1996; Division of Communicable Disease Control, Office of Statistics and Surveillance; Office of AIDS, AIDS Case Registry; Genetic Disease Branch, Newborn Screening Program. Department of Finance, State Census Data Center: 1997 Race/Ethnic Population by County with Age and sex detail, June 1999.